

ins
C1

Porcine Retrovirus

The present invention relates inter alia to porcine retrovirus (PoEV) fragments, in particular polynucleotide fragments encoding at least one porcine retrovirus expression product, a recombinant vector comprising at least one polynucleotide fragment, use of PoEV polynucleotide fragments in the detection of native porcine retrovirus, a host cell containing at least one PoEV polynucleotide fragment or a recombinant vector comprising at least one PoEV polynucleotide fragment, PoEV polypeptides, antibodies immuno-reactive with PoEV polypeptides, pharmaceutical compositions comprising recombinant PoEV polypeptides for use as prophylactic and/or therapeutic agents and uses of PoEV polynucleotide fragments and/or polypeptides in medicine, including veterinary medicine and in the preparation of medicaments for use in medicine, including veterinary medicine.

Porcine retrovirus (PoEV) is an endogenous (genetically acquired) retrovirus isolated from pigs and expressed in cell lines derived from porcine material. There are no known pathogenic effects associated with the virus per se in its natural host although the virus appears to be associated with lymphomas in pigs and related viruses are associated with leukaemias and lymphomas in other species. The virus has been reported to infect cells from a variety of non-porcine origins and is, therefore, designated as a xenotropic, amphotropic or polytrophic virus (Lieber MM, Sherr CJ. Benveniste RE and Todaro

GJ. 1975; Strandstrom H, Verjalainen P, Moening V, Hunsmann G, Schwarz H, and Schafer W. 1974; Todaro GJ, Benveniste RE, Lieber MM and Sherr CJ. 1974). The observation that the above viruses may have the potential to infect humans and have a pathogenic effect suggests that the issue of porcine retroviruses must be addressed in the context of xenotransplanting pig tissues or cells. Therefore, information on the properties of PoEV and the development of diagnostic reagents, molecular engineering tools and potential vaccine materials is of paramount importance for example in xenotransplantation technology and the like.

It is an object of the present invention to obviate and/or mitigate against at least some of the above disadvantages.

In one aspect the present invention provides an isolated PoEV polynucleotide fragment:

- (a) encoding at least one porcine retrovirus (PoEV) expression product;
- (b) encoding a physiologically active and/or immunogenic derivative of said expression product; or
- (c) which is complementary to a polynucleotide sequence as defined in (a) or (b).

Preferably, the polynucleotide fragment encodes the gag gene (gag), polymerase gene (pol) and/or envelope (env) gene of PoEV. Thus, said expression product can be the virion core polypeptides (GAG) and polymerase (POL) and/or envelope (ENV) polypeptides of PoEV. Thus, the invention further provides a recombinant PoEV virion core, polymerase and/or envelope polypeptide.

"Polynucleotide fragment" as used herein refers to a chain of nucleotides such as deoxyribose nucleic acid (DNA) and

603020 "EST 2000"

ins
C2

transcription products thereof, such as RNA. Naturally, the skilled addressee will appreciate the whole naturally occurring PoEV genome is not included in the definition of polynucleotide fragment.

The polynucleotide fragment can be isolated in the sense that it is substantially free of biological material with which the whole genome is normally associated *in vivo*. The isolated polynucleotide fragment may be cloned to provide a recombinant molecule comprising the polynucleotide fragment. Thus, "polynucleotide fragment" includes double and single stranded DNA, RNA and polynucleotide sequences derived therefrom, for example, subsequences of said fragment and which are of any desirable length. Where a nucleic acid is single stranded then both a given strand and a sequence complementary thereto is within the scope of the present invention.

In general, the term "expression product" refers to both transcription and translation products of said polynucleotide fragments. When the expression product is a "polypeptide" (i.e. a chain or sequence of amino acids displaying a biological and/or immunological activity substantially similar to the biological and/or immunological activity of PoEV virion core, polymerase and/or envelope protein), it does not refer to a specific length of the product as such. Thus, the skilled addressee will appreciate that "polypeptide" encompasses *inter alia* peptides, polypeptides and proteins of PoEV. The polypeptide if required, can be modified *in vivo* and *in vitro*, for example by glycosylation, amidation, carboxylation, phosphorylation and/or post-translational cleavage.

Polynucleotide fragments comprising portions encompassing the PoEV genome, and derived from retrovirus particles released from a reverse transcriptase-positive porcine kidney cell line PK-15, have been molecularly cloned into a plasmid vector. This was achieved by synthesising cDNAs of PoEV RNA genomes which were recovered from porcine kidney cells expressing the endogenous virus. The cDNA was cloned into a plasmid vector and the isolated PoEV DNA fragment determined (see Figures 1,2 and 3). The sequence of the sequence identified in Figure 1 was the earliest determined sequence, followed by the sequence in Figure 2 and lastly by the most recently revised sequence shown in Figure 3. An additional study has been carried out to determine whether or not the human cell line "Raji" was susceptible to infection with the PoEV present in porcine kidney cells (PK15). A raji clone has now been obtained and the DNA sequence of its env gene region has been determined (see Figure 4).

The DNA fragment of Figure 3 was shown to encode three open reading frames (ORFs) of 524, 1194 and 656 amino acids respectively.

A comparison of the amino acid sequence against previously sequenced retroviruses from other species indicated that novel retrovirus cDNA had been cloned. Sequence analysis using the Lasergene software from DNASTAR Inc. showed that homologies were observed between the cloned PoEV DNA and the majority of retroviruses and that the closest homologies were to gibbon leukaemia virus (GaLV) in the polymerase (pol) and (env) regions of the pro-virus.

The first open reading frame ORF of Figure 3 (nucleotides 588-

2162) is predicted to encode the PoEV virion core polypeptide (gag gene). The second ORF (nucleotides 2163-5747) is predicted to encode the PoEV polymerase polypeptide (pol gene). The third ORF (nucleotides 5620-7590) is predicted to encode the PoEV envelope polypeptide (env gene). The skilled addressee will appreciate that it is possible to genetically manipulate the polynucleotide fragment or derivatives thereof, for example to clone the gene by recombinant DNA techniques generally known in the art and to express the polypeptides encoded thereby in vitro and/or in vivo. DNA fragments having the polynucleotide sequence depicted in Figures 1, 2, 3 and/or 4 or DNA/RNA derivatives thereof, may be used as a diagnostic tool or as a reagent for detecting PoEV nucleic acid in nucleic acids from donor animals or as a vaccine.

Preferred fragments of this aspect of the invention are polynucleotide fragments encoding: (a) at least one of the one to three polypeptides having an amino acid sequence which is shown in Figures 1,2,3 and/or 4 (b) encoding a polypeptide which is a physiologically active and/or immunogenic derivative of at least one of the polypeptides defined in (a); or (c) which is complementary to a polynucleotide sequence as defined above; or polynucleotide fragments: (a) comprising at least one of the ORFs shown in Figures 1,2,3 and/or 4 or comprising a corresponding RNA sequence; (b) comprising a sequence having substantial nucleotide sequence identity with a sequence as described in (a) above; or (c) comprising a sequence which is complementary to a sequence as described in (a) or (b) above. It is to be understood that the term "substantial sequence identity" is taken to mean at

1



Variable	Mean	SD	Min	Max	Skewness	Kurtosis	Normality
Age	38.5	12.5	25	65	0.1	3.2	0.95
Gender	0.5	0.5	0	1	0.0	3.0	0.98
Marital Status	0.7	0.5	0	1	0.1	3.1	0.96
Education	12.5	2.5	9	16	0.2	3.3	0.94
Income	1500	500	500	3000	0.3	3.4	0.92
Health	0.8	0.4	0	1	0.2	3.2	0.95
Stress	4.5	1.5	1	7	0.4	3.5	0.90
Depression	3.2	1.2	1	5	0.3	3.4	0.91
Life Satisfaction	5.8	1.0	4	7	-0.1	3.1	0.97
Resilience	6.5	1.5	4	8	-0.2	3.2	0.96
Optimism	6.2	1.2	4	8	-0.1	3.1	0.97
Self-Esteem	6.0	1.0	4	8	-0.1	3.1	0.97
Emotional Stability	6.8	1.0	4	8	-0.2	3.2	0.96
Life Satisfaction	5.8	1.0	4	7	-0.1	3.1	0.97
Resilience	6.5	1.5	4	8	-0.2	3.2	0.96
Optimism	6.2	1.2	4	8	-0.1	3.1	0.97
Self-Esteem	6.0	1.0	4	8	-0.1	3.1	0.97
Emotional Stability	6.8	1.0	4	8	-0.2	3.2	0.96

If a PoEV specific test polynucleotide sequence is to be used in hybridisation studies, to test for the presence of PoEV nucleic acid in a sample, the test polynucleotide should preferably remain hybridised to a sample polynucleotide under

stringent conditions. If desired, either the test or sample polynucleotide may be immobilised. Generally the test polynucleotide sequence is at least 10 or at least 50 bases in length. It may be labelled by suitable techniques known in the art. Preferably the test polynucleotide sequence is at least 200 bases in length and may even be several kilobases in length. Thus, either a denatured sample or test sequence can be first bound to a support. Hybridization can be effected at a temperature of between 50 and 70°C in double strength SSC (2xNaCl 17.5g/l and sodium citrate (SC) at 8.8g/l) buffered saline containing 0.1% sodium dodecyl sulphate (SDS). This can be followed by rinsing of the support at the same temperature but with a buffer having a reduced SSC concentration. Depending upon the degree of stringency required, and thus the degree of similarity of the sequences, such reduced concentration buffers are typically single strength SSC containing 0.1%SDS, half strength SSC containing 0.1%SDS and one tenth strength SSC containing 0.1%SDS. Sequences having the greatest degree of similarity are those the hybridisation of which is least affected by washing in buffers of reduced concentration. It is most preferred that the sample and inventive sequences are so similar that the hybridisation between them is substantially unaffected by washing or incubation in one tenth strength sodium citrate buffer containing 0.1%SDS.

PoEV specific oligonucleotides may be designed to specifically hybridise to PoEV nucleic acid. They may be synthesised, by known techniques and used as primers in PCR or sequencing reactions or as probes in hybridisations designed to

00174553-000000

the DNA is PoEV.

The present invention also encompasses PoEV detection kits including at least one oligonucleotide which is PoEV specific, as well as any necessary reaction reagents, washing reagents, detection reagents, signal producing agents and the like for use in the test formats outlined above.

In a further aspect there is also provided use of a PoEV specific polynucleotide in the detection of PoEV in a sample.

In a yet further aspect there is provided use of a PoEV specific polynucleotide in a PCR for the detection of PoEV in a sample.

The skilled addressee will appreciate how polynucleotide fragments may be designed and used as primers/probes in polymerase chain reaction (PCR) experiments or Southern analysis (i.e. hybridisation studies) for detecting the presence or otherwise of PoEV polynucleotide in the nucleic acid of pigs or in cell, tissue or organ samples taken from pigs (e.g. from potential transplant organs such as liver, kidney and heart). Such cells, tissues or organs can be derived from transgenic animals produced as described in EP-A-0493852, or by other means known in the art. Thus the cells, tissues or organs of transgenic pigs can be associated with one or more homologous complement restriction factors active in humans to prevent/reduce activation of complement.

Furthermore the polynucleotide fragments of the present invention can be used to analyze the genetic organisation of endogenous PoEV located in the animal cell genome in pigs thus permitting the screening of herds of pigs for altered provirus

and genomic loci (e.g. non-expressed provirus loci). Such a screening method would facilitate, for example, screening in a population of animals which are bred to lack expressed provirus and genomic loci and/or loci that do not encode infectious virus particles.

Reagents may also be developed from said polynucleotide fragments as aids to develop pigs that do not express an infectious, PoEV capable of infecting humans. Such pigs could still contain partial defective genomes that could result in the expression of non-infectious particles, viral proteins or viral mRNA. Alternatively, it may be possible to use constructs derived from the PoEV polynucleotide sequence to act as insertional mutagens to knockout the productive infectious PoEV in embryos, embryonic stem cells, or cells containing totipotent nuclei capable of forming a viable embryo. Thus gag, pol and/or env gene "knockouts" may be constructed to allow development of breeding programmes in pigs whereby endogenous PoEV is substantially prevented or reduced. For example the nucleotide sequence of PoEV can be manipulated e.g. by deletion of a coding sequence in vitro and the resulting construct used to replace the natural PoEV sequence by recombination. Thus, the proviral genome can be rendered inactive in the porcine cells. The knockouts can be manipulated into embryos and/or stem cells and if required manipulated nuclei can be transferred from target cells to germ cells using micromanipulation techniques well known in the art. The invention also extends to animals derived from such germ cells.

Thus, transgenic pigs may be produced containing anti-sense

constructs and/or ribozyme constructs capable of downregulating the expression of viral proteins, or transgenic pigs expressing a single chain immunoglobulin molecule with specificity for PoEV proteins or other protein that might interfere with protein synthesis or viral assembly may also be produced. Similar transgenes encoding trans-dominant negative regulators of PoEV expression or transgenes encoding competitive defective "genomic RNAs" may be used to reduce or eliminate the production of infectious virions. The generation of reagents to suppress the expression of native PoEV loci in pigs, such as, by generation of antisense nucleic acids (e.g. antisense mRNAs), ribozymes or other antiviral reagents may also be developed.

The polynucleotide fragment can be molecularly cloned into a prokaryotic or eukaryotic expression vector using standard techniques and administered to a host. The expression vector is taken up by cells and the polynucleotide fragment of interest expressed, producing protein. Presentation of the protein on cell surface stimulates the host immune system to produce antibodies immunoreactive with said protein as part of a defence mechanism. Thus, expressed protein may be used as a vaccine.

Inactivated vaccines can be produced from PoEV's or cells releasing PoEV. Such infected cells may be generated by natural infection or by transfection of a proviral clone of PoEV. It will be understood that a proviral clone is a molecular clone encoding on at least one antigenic polypeptide of PoEV. After harvesting the virus and/or the infected cells, viruses or infected cells present can be inactivated for example, with formaldehyde, gluteraldehyde, acetylenimine or other

Proviral clones of PoEV can be engineered to develop single cycle or replication defective viral vectors suitable for vaccination using techniques. Such viral vectors known in the art (e.g. MuLV Murine Leukaemia Retrovirus, Adenovirus and Herpesviruses (Anderson WF. (1992). Human Gene Therapy. Science 256, 808-813) may have one or more genes essential for replication deleted, with the missing gene function expressed constitutively or conditionally from a further, different construct which is integrated into the chromosomal DNA of a complementing cell line to the proviral PoEV clone. PoEV virions released from the cell line may infect secondary target cells in the vaccinee but not produce further infectious virus particles. For instance, the polynucleotide sequence encoding the reverse transcriptase domain of *pol* can be deleted from the proviral PoEV clone and the reverse transcriptase domain of *pol* integrated into the complementing cell line.

It will be understood that the polynucleotides; polypeptides; PoEV free cells, tissues and/or organs encompassed

by the present invention could be used in therapy, diagnosis, and/or methods of treatment. The polynucleotides; polypeptides; PoEV free cells, tissues and/or organs encompassed by the present invention can also be used in the preparation of medicaments for use in therapy or diagnosis.

The cloning and expression of a recombinant PoEV polynucleotide fragment also facilitates in producing anti-PoEV antibodies and fragments thereof (particularly monoclonal antibodies) and evaluation of in vitro and in vivo biological activity of recombinant PoEV polymerase and/or envelope polypeptides. The antibodies may be employed in diagnostic tests for native PoEV virus.

It will be understood that for the particular PoEV polypeptides embraced herein, natural variations can exist between individuals or between members of the family Suidae (i.e. the pig family). These variations may be demonstrated by (an) amino acid difference(s) in the overall sequence or by deletions, substitutions, insertions, inversions or additions of (an) amino acid(s) in said sequence. All such derivatives showing active polymerase and/or envelope polypeptide physiological and/or immunological activity are included within the scope of the invention. For example, for the purpose of the present invention conservative replacements may be made between amino acids within the following groups:

- (I) Alanine, serine, threonine;
- (II) Glutamic acid and aspartic acid;
- (III) Arginine and leucine;
- (IV) Asparagine and glutamine;

09171553-020099

(V) Isoleucine, leucine and valine;

(VI) Phenylalanine, tyrosine and tryptophan

Moreover, recombinant DNA technology may be used to prepare nucleic acid sequences encoding the various derivatives outlined above.

As is well known in the art, the degeneracy of the genetic code permits substitution of bases in a codon resulting in a different codon which is still capable of coding for the same amino acid, e.g. the codon for amino acid glutamic acid is both GAT and GAA. Consequently, it is clear that for the expression of polypeptides with the amino acid sequences shown in Figure 1 or fragments thereof, use can be made of a derivative nucleic acid sequence with such an alternative codon composition different from the nucleic acid sequence shown in said Figure 1.

Furthermore, fragments derived from the PoEV core, polymerase and/or envelope polypeptides as depicted in Figure 3, which still display PoEV virus core polypeptide, polymerase and/or envelope polypeptide properties, or fragments derived from the nucleic acid sequence encoding the virus core polypeptides, polymerase and/or envelope polypeptides or derived from the nucleotide sequence depicted in Figures 1, 2, 3 and/or 4 encoding fragments of said virus core polypeptide, polymerase and/or envelope polypeptides are also included of the present invention. Naturally, the skilled addressee will appreciate within the ambit that the said fragments should substantially retain the physiological and/or immunological properties of the GAG, POL and/or ENV polypeptides.

The PoEV polynucleotide fragment of the present invention

A polynucleotide fragment according to the present invention can be ligated to various expression controlling sequences, resulting in a so called recombinant nucleic acid molecule. Thus, the present invention also includes an expression vector containing an expressible PoEV nucleic acid molecule. The recombinant PoEV nucleic acid molecule can then be used for the transformation of a suitable host. Such hybrid molecules are preferably derived from, for example, plasmids or from nucleic acid sequences present in bacteriophages or viruses and are termed vector molecules.

The methods to be used for the construction of a recombinant nucleic acid molecule according to the invention are known to those of ordinary skill in the art and are *inter alia* set forth in Sambrook, et al. (Molecular Cloning: a laboratory manual Cold Spring Harbour Laboratory, 1989).

The present invention also relates to a transformed cell containing the PoEV polynucleotide fragment in an expressible form. "Transformation", as used herein, refers to the introduction of a heterologous polynucleotide fragment into a

—

Examples Section

Example 1

Preparation of viral RNA

500ml of supernatant derived from exponentially growing porcine kidney cells (PK-15, American Type Culture Collection CCL 33) was clarified by centrifugation of approximately 11,000xg for 10 minutes. Virus was pelleted from the clarified supernatant by centrifugation at approximately 100,000xg for 60 minutes. The supernatant was discarded and the viral pellet retained for the preparation of viral RNA genomes. RNA was prepared from the virus pellet using a Dynabeads (registered trade mark) mRNA Direct kit according to the manufacturer's protocols; A PoEV virus pellet was resuspended in 500 μ l of TNE (10mM Tris HCl pH8.0, 0.1M NaCl, 1mM EDTA) and the virions disrupted by the addition of 2ml of lysis/binding buffer. Dynabeads Oligo(dT)₂₅ were conditioned according to the manufacturer's instructions and added to the virus disrupted solution. Viral RNA was allowed to bind to the Dynabead for 10 minutes before the supernatant was removed and the bound RNA was washed three times with washing buffer with LiDS (0.5ml) and twice with washing buffer alone. The RNA was finally resuspended in 25 μ l of elution solution. All procedures were performed at ambient temperature. RNase contamination was avoided by the wearing of gloves, observation of sterile technique and treatment of solutions and non-disposable glass and plasticware with diethyl pyrocarbonate (DEPC). The RNA was resuspended in DEPC- treated sterile water.

Example 2

Synthesis of cDNA

cDNA was synthesised from the purified genomic RNA using Great Lengths™ cDNA amplification reverse transcriptase reagents (Clontech Laboratories Inc.) following the manufacturer's instructions. The RNA was primed with both oligo(dT) and random hexamers to maximise synthesis.

The Great Lengths cDNA synthesis protocol is based on a modified Gubler and Hoffman (1983) protocol for generating complementary DNA libraries and essentially consists of first-strand synthesis, second strand synthesis, adaptor ligation, and size fractionation.

First strand synthesis: lock-docking primers anneal to the beginning of the poly-A tail of the RNA due to the presence of A, C or a residue at the 3'-end of the primer. This increases the efficiency of cDNA synthesis of eliminating unnecessary reverse transcription of long stretches of poly-A. In addition, the reverse transcriptase used is MMLV (RNase H⁻) which gives consistently better yields than do wild-type MMLV or AMV reverse transcriptase.

Second strand synthesis: the ratio of DNA polymerase I for RNase H has been optimised to increase the efficiency of the second strand synthesis and to minimize priming by hair pin loop formation. Following second-strand synthesis, the ds cDNA is treated with T4 DNA polymerase to create blunt ends.

Adaptor ligation: the cDNA is ligated to a specially designed adaptor that has a pre-existing EcoRI "sticky end". The use of this adaptor, instead of a linker, eliminates the need to methylate and the EcoRI - digest the cDNA, and thus leaves internal EcoRI, sites intact. The adaptor is 5'-phosphorylated at the blunt end to allow efficient ligation to the blunt-ended cDNA.

Size fractionation: the ds cDNA is phosphorylated at the EcoRI sites and size-fractionated to remove unligated adaptors and unincorporated nucleotides. The resulting cDNA is ready for cloning into a suitable EcoRI-digested vector.

Example 3

Molecular cloning of cDNA

The size fractionated fragment was ligated with EcoR I- digested pZERO™ -1 plasmid vector DNA (Invitrogen Corporation, San Diego, U.S.). The ligation mix was used to transform competent TOP10F' cells and these were plated onto L-Agar containing zeocin following the manufacturer's instructions (Zero Background™ cloning kit - Invitrogen). Several of the resulting zeocin resistant colonies were amplified in L-Broth containing zeocin and the plasmid DNA was purified by alkaline lysis (Maniatis et al., 1982).

The plasmid DNA was digested to completion with the endonuclease EcoR I and the resulting DNA fragments were separated by electrophoresis through an 1.0% agarose gel (Maniatis et al., 1982), in order to check that a fragment in the

predicted size fractionated size range had been cloned. A clone identified as pPoEV was used in further experimentation.

Example 4

DNA sequence analysis.

pPoEV plasmid DNA was purified according to common techniques (Sambrook et al, 1989) and sequenced using an ABI automated sequencer. Overlapping sequencing primers from both strands of the molecular clone were used to determine the nucleotide sequence.

The first sequence obtained is shown in Figure 1. This sequence was identified as encoding two ORFs of 924 (nucleotides 23-2793) and 213 (nucleotides 2642-3297) amino acids, relating to the *pol* and *env* genes respectively. This sequence was revised and updated to the second sequence as shown in Figure 2. This second sequence was identified as encoding three ORFs of 516 (nucleotides 576-2126), 1136 (nucleotides 2143-5733) and 656 (nucleotides 5606-7576) amino acids, encoding the PoEV *gag*, *pol* and *env* genes respectively. This second sequence has since been revised and updated to the sequence shown in Figure 3. This third sequence was identified as encoding three ORFs of 524 (nucleotides 588-2162), 1194 (nucleotides 2163-5747) and 656 (nucleotides 5620-7590) amino acids, encoding the PoEV *gag*, *pol* and *env* genes respectively.

The differences in the disclosed sequences is reflected by improvements in carrying out and analysing the sequence obtained. However, there is 100% identity at the nucleic acid level, between positions 21-2681 of the first sequence and positions 2972-5653 of the third sequence. Overall there is a 70.5%

identity in the entire 3310 bp of the first sequence with a corresponding portion of the third sequence.

There are only 3 base changes between the second sequence and the third sequence. These are as follows:

<u>base no. (from Figure 2)</u>	<u>change</u>
2121	insertion of a "G"
2157	insertion of a "G"
5902	"R" to an "A"
7700	"M" to an "A"

ins C3
~~The changes at base nos. 5902 and 7700 do not effect the corresponding amino acid sequence. However, the changes at positions 2121 and 2157 alter the amino acid sequence at the end of GAG and the begining of POL. For GAG the final amino acid "S" have now been replaced by "VLAL EEDKD". The total product size is now 524 amino acids. For POL, the first five amino acids "RLGET" have been deleted and replaced by "GRR". The total product size is now 1194 amino acids.~~

Similarities were observed between pPoEV and the majority of retroviruses determined by using alogrithims from DNASTAR Inc. Lasergene software (DNASTAR). The similarities were closest with gibbon ape leukaemia virus (GaLV) in the polymerase (pol) regions of the pro-virus at 63.5%, in the virus core (gag) region, 59.2% and in the envelope (env) region, 39.3% The nucleotide sequence and major ORFs of the pPoEV insert are shown in Figure 3. The largest ORF (nucleotides 2163-5747) encodes the polymerase polypeptide and the smaller ORFs (nucleotides 538-2162 and 5620-7590) encode the core and envelope polypeptides respectively.

Example 5

Purification of cellular DNA from cultured cells, tissues and blood.

Cultured cells

Cells were maintained in culture and approximately 5×10^7 cells were harvested for DNA preparation. The cells were pelleted by centrifugation resuspended in phosphate-buffered saline, re-centrifuged at 1000g for 2 minutes and the supernatant was discarded.

Porcine tissues

Porcine tissue samples were frozen in liquid nitrogen and powdered by grinding in a mortar or between metal foil. The samples were resuspended in 5ml of extraction buffer consisting of 0.025M EDTA (pH 8.0), 0.01M Tris.Cl pH 8.0, 0.5% SDS 20 μ g/ml RNase and 100 μ g/ml proteinase K (Maniatis et al., 1982).

Porcine blood

A buffy coat was prepared from the blood samples. 20ml samples were centrifuged at 1000g for 15 minutes. The buffy coat was resuspended in buffer and the samples centrifuged at 1000g for 15 minutes. The process was repeated one further time. The sample was mixed with 5ml (3x volume) of extraction buffer (Maniatis et al., 1982).

Purification

The samples (i.e. cultured cells, porcine tissue or porcine blood cells) in proteinase K-extraction buffer containing 20 μ g/ml RNase

66000-655460

Example 6

In order to demonstrate that the molecularly cloned DNA comprising the insert from PoEV was derived from the PK-15 cell line (American Type Culture Collection CCL33), the DNA was hybridised against cellular DNAs and its ability to detect proviral DNA was examined.

The DNAs probed were as follows :

- a) Copy number controls of pPoEV DNA linearized by digestion with EcoRI. One copy per haploid cell genome was estimated to be 6.84pg. The control was present at an estimated copy number of 1, 5 and 10 copies.
- b) PK-15 DNA.
- c) Negative control HeLa (American Type Culture Collection

CCL2) DNA derived from a human adenocarcinoma cell line harbouring human papillomavirus type 18 DNA.

- d) Negative control SP20 (European Collection of Animal Cell Cultures 85072401) DNA derived from a murine myeloma cell line harbouring a xenotropic MuLV retrovirus.

A hybridisation signal was observed in only the PK-15 porcine DNA. No signal was detected in either the negative human or murine DNAs. The PK-15 DNA contained more than 10 copies per cell with an estimated copy number of 20. The sizes of the three major EcoRI- endonuclease digested DNA fragments were approximately 3.8kb, 1.8kb and 0.6kb. The sizes of relevant fragments detected in the recombinant pPoEV were comparable.

There are, as expected, a number of fragments common to the genomic DNA of PK-15 and pPoEV DNA and there is agreement between the patterns observed in both DNAs digested with XhoI, BamHI and HindIII. However, there are additional fragments obtained on digestion of pPoEV DNA by a number of endonucleases.

pPoEV sequences were also detected in swine testes (American Type Culture Collection CRL 1746) and primary porcine kidney cells (Central Veterinary Laboratory batch C04495) but not in hamster CHOK1 (American Type Culture Collection CCL61) or murine NS0 myeloma cells (European Collection of Animal Cell Cultures 85110503).

09455-02099
000020-0554260

In order to demonstrate that the molecularly cloned DNA comprising the insert from pPoEV could detect sequences in porcine cells and tissues in addition to PK-15 the pPoEV DNA was hybridised against cellular DNA from tissues derived from pigs and its ability to detect proviral DNA was examined (Maniatis et al., 1982).

The DNA purified from pPoEV was radioactively labelled and used to probe a Southern blot of endonuclease digested DNAs derived from pig organs including liver, kidney, heart and blood.

The DNAs probed were as follows :

- a) Copy number controls of pPoEV DNA linearized by digestion with EcoRI. One copy per haploid cell genome was estimated to be 6.84pg. The control was present at an estimated copy number of 5, 10, 20 and 50 copies.
- b) DNA purified from the porcine tissues digested with EcoRI.

A hybridisation signal was observed in all the porcine DNAs.

The DNAs contained less than 5 copies per cell. There were approximately eight distinct bands in each DNA. The sizes of the three major endonuclease digested DNA fragments were approximately 3.8kb, 1.8kb and 0.6kb.

Example 7

Polymerase Chain Reaction (PCR) Amplifications

~~Oligonucleotides were selected from the PoEV genome.~~

The upstream primer was 5'-GGA AGT GGA CTT CAC TGA G-3'.

~~The downstream primer was 5'-CTT TCC ACC CCG AAT CCG -3'.~~

The PCR was performed as described by Saiki et al (1987). One $1\mu\text{l}$ of $100\text{ng}/\mu\text{l}$ template DNA was added to a $49\mu\text{l}$ reaction mixture containing $200\mu\text{M}$ of dATP, dCTP, dGTP, dTTP, 30pmol of both primers from the pair described above, 1 unit of DNA polymerase and $5\mu\text{l}$ of reaction buffer. The reaction buffer contained 200mM Tris-HCl pH 8.4, 500mM potassium chloride and 15mM magnesium chloride, ultrapure water. The solution was overlaid with two drops of mineral oil to prevent evaporation. Thirty five cycles of amplification were performed using a Perkin Elmer Cetus thermal cycler. Each cycle consisted of 1 minute. at 95°C to denature the DNA, 1 minute. at 53°C to anneal the primers to the template and 1 minute. at 72°C for primer extension. After the last cycle a further incubation for 10 minutes. at 72°C was performed to allow extension of any partially completed product. On completion of the amplification, $10\mu\text{l}$ of the reaction mixture was electrophoresed through a 5 per cent acrylamide gel. The DNA was visualised by staining with ethidium bromide and exposure to ultraviolet light (320nm).

The PCR reaction amplified a sequence of approximately 787bp from pPoEV and from porcine cells as expected indicating that the assay detected the PoEV proviral DNA. There was no specific amplification of the expected sequence in cells of non-porcine origin and therefore, the PCR reaction and recombinant clone can be used as a specific and sensitive diagnostic tool for detection of PoEV.

~~Two further digonucleotides were designed against the 3' end of the pol gene and 5' end of the gag gene respectively.~~

~~The 3' pol oligionucleotide was 5'-GAT GGC TCT CCT GCC CTT TG-3'~~

~~The 5' gag oligionucleotide was 5'-CGA TGG AGG CGA ACC TTA AGG-3'~~

The above oligionucleotide were also used in in PCR reactions according to the conditions described above, with the exceptions that the annealing temperature was 58° and 30 cycles of replication were carried out. The PCR reaction amplified a sequence of approximately 468bp from pPoEV and from porcine cells.

Example 8

Production of PoEV polypeptide in *Escherichia coli*.

The open reading frame (ORF) encoding the pol peptide was isolated from the pPoEV clone and molecularly cloned into the plasmid pGEX-4T-1 (Pharmacia Ltd.) for expression.

Two ml cultures of *E. coli* transformed with various expression constructs were grown with shaking at 37°C to late log phase

(O.D._{600nm} of 0.6) and induced by the addition of IPTG to 0.1 mM. Induced cultures were then incubated for a further 2 hours after which the bacteria were collected by centrifugation. The bacterial pellet was lysed by boiling in SDS-PAGE sample buffer and the protein profile of the induced bacteria was analysed on a 12% acrylamide gel (Laemmli, 1970) followed by staining with coomassie brilliant blue dye.

Example 9

Isolation and partial sequencing of Raji clone

The aim of the study was to determine whether the human cell line "Raji" was susceptible to infection with the PoEV present in porcine kidney cells (PK15). In order to test the capacity of the virus for xenotropism, PK15 cells were co-cultured with the B lymphoblastoid (Raji) cell line over 20 passages.

The culture system utilised direct culture and transwells, which separated the human and porcine cells, but permitted viruses to pass through the separating membrane. After every fifth passage, supernatants from the human cell lines are tested for the presence of retrovirus by reverse transcriptase assay.

Cell cultures

Porcine kidney (PK15) cells (ATCC CCL 33) were used as the source of PoEV. The human cells used for co-cultivation with PK15 cells were the lymphoblast-like Burkitts lymphoma Raji (ATCC CCL 86) cell line. This cell line does not harbour endogenous

retroviruses and lacks reverse transcriptase activity when tested by the present inventors.

Co-cultivation

Raji cells were co-cultivated directly with PK15 cells in duplicate 80cm² flasks and exposed to the PK15 cells throughout the 20 passage culture period. The cells were passaged twice per week and PK15 cells added as necessary from a stock culture. At every fifth passage a sample of Raji cells was removed from the co-culture, washed and cultured for 3-4 days. Supernatant was then harvested and tested for presence of retrovirus by reverse transcriptase (RT) assay.

RESULTS

The presence of reverse transcriptase activity with a preference for the Mn²⁺ cation in the supernatant from detector cell cultures is suggestive of infection by porcine retrovirus. Reverse transcriptase activity with preference for the Mn²⁺ template was not detected in the duplicate co-cultivated test cultures at passage 5 but was detected at passages 10, 15 and 20. No significant RT activity was detected in the negative control cultures. RT activity with preference for the Mn²⁺ template was detected in positive control cultures at passage 5 and 20.

An infected raji culture was diluted to single cells, and then a selection of cells cultured separately such that each culture originated from one cell. Each culture was tested by reverse-transcriptase assay.

Genomic DNA was made from an RT-positive clone as described in example 5 -purification. The PoEV ENV region was amplified by PCR as described below and the product molecularly cloned into pMOS blue T-vector (Amersham). This molecular clone was then sequenced (Fig. 4).

PCR

~~Oligonucleotides were selected from the PoEV genome.~~

The upstream primer was 5'-GAT GGC TCT CCT GCC CTT TG -3'

5' base position: 5240

The downstream primer was 5'-CCA CAG TCG TAC ACC ACG -3'

5' base position: 8144

~~Expected product size: 2904bp~~

Approx. 1 μ g of genomic raji clone DNA was added to a 50 μ l reaction mixture containing 200 μ M of dATP, dCTP, dGTP, dTTP, 30pM each primer detailed above, 1u Taq DNA polymerase and 5 μ l reaction buffer. The reaction buffer contained 200mM Tris.Cl pH 8.4, 500mM potassium chloride, 15mM magnesium chloride and ultrapure water. The solution was overlaid with two drops of mineral oil to prevent evaporation. Thirty cycles of amplification was performed followed by an elongated extension reaction of 60min. at 72°C.

The cycles consisted of:

95°C 1 min.

56°C 1 min.

72°C 2 min.

The PCR product was visualised as described in example 7.
Product size: ~3Kb.

CLONING

The PCR product was molecularly cloned into pMOS-Blue T-vector as directed by the manufacturer (pMOS-Blue T-vector kit - Amersham).

20 transformed colonies (clones) were picked and added to 5mls L-broth containing 50 µg/ml ampicillin. The cultures were grown shaking at 37°C overnight. Plasmid DNA was isolated from each clone using the perfect prep plasmid isolation kit as directed by the manufacturer (5 Prime - 3 Prime Inc. Boulder, CO, USA).

Plasmid DNA was digested to completion with the endonucleases EcoRI and HindIII and the products visualised on an ethidium bromide-stained 1% agarose gel. A clone (raji env clone) showing the same banding pattern as that predicted for 'PK15 cell line derived PoEV', was selected for sequencing.

SEQUENCING

Raji env clone plasmid DNA prepared above was sequenced using an ABI automated sequencer, and the commercially available T7 sequencing primer.

Example 10

Phylogenetic analysis was performed using the PHYLIP package. Sequence distances were calculated using the PROTDIST program (Dayhoff matrix) and a neighbour-joining unrooted phylogenetic tree reconstructed using the NEIGHBOUR program. Bootstrapping was performed using 200 replicates of the pol alignment, created using the SEQBOOT program and a consensus tree was obtained using the CONSENSE program (see Figure J). The bootstrap percentages are indicated at the branch fork, with missing values equal to 100%. The data indicate that PoEV is closely related to but distinct from the type-C oncovirus typified by gibbon, murine and feline leukaemia viruses. A phylogenetic tree was constructed from the pol alignment using the maximum likelihood algorithm (Dayhoff matrix). This tree differed from the pol NJ tree only in the placement of the BaEV lineage in relation to other mammalian type C viruses and corresponded to a low bootstrap for the BaEV fork observed in the NJ tree.

Example 11Analvsis of the LTR and adjacent region

The long terminal repeat (LTR) is a reiterated sequence at each end of the provirus that contains the enhancer and promoter governing transcription of the provirus as well as sequences required for reverse transcription of the RNA genome and integration of the proviral DNA. Three recognised domains of the LTR are identifiable, U3, R and U5 with the LTR being delineated by inverse repeats AATGAAAGG and CCTTTCATT at the 5' and 3' ends of U3 and U5 respectively.

<u>LTR Domain</u>	<u>PoEV Genome Sequence</u>	<u>Length bp</u>
in accordance with Figure 3		
U3	7638-8106	469
R*	8107-8138, 1-61	82
U5	62-143	82

*The position of the R is defined here by similarity to the 3' end of the MuLV LTR and is compatible with the observed location of a cap site approximately 24 bp downstream of the TATA box.

The U3 region contains multiple potential transcription sites as shown in Figure 6. Most of the U3 region shows little or no homology to other mammalian type-C retroviruses which show conserved sites or repeat elements. However, there is homology to other mammaliann type-C viruses towards the 3' end of the U3 & region and into R and U5. Amongst the potential transcription factor sites are those for the following:

E47 is the prototype member of a new family of tissue specific enhancer proteins that have been shown to bind to the enhancer of murine leukaemia virus.

ETS-1 is a transcription factor primarily expressed in the haematopoietic lineage.

The LTR contains direct repeats at 30006-3062 and 3045-3101 which together contain three potential CCAATT boxes. A potential TATA box is located at position 3129-3144.

The R region contains a PADS (Poly A downstream element) and consensus polyadenylation signal (AATAAA).

The primer binding site (PBS) of PoEV is glycine(2) tRNA which has not reported for any exogenous retrovirus.

Clare JJ, Rayment FB, Ballantine SP, Sreekrishna K and Romanos MA. (1991). High Level expression of tetanus toxin fragment C in *Pichia pastoris* strains containing multiple tandem integrations of the gene. *Biototechnology*, 9, 455-460

Derynck R, Singh A and Goeddel DV. (1983). Expression of the human interferon- γ in yeast. *Nucleic Acids Res.*, 11, 1819-1837.

DNASTAR. (1994). Lasergene Biocomputing Software for Windows. User's Guide.

Invitrogen. Version A. Zero Background™ Cloning Kit Catalog no K2500-01.

Laemmli UK. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4N. *Nature*, 227, 680-685.

Lieber MM, Sherr CJ, Benveniste RE and Torado GJ. (1975). Biologic and immunologic properties of porcine type C viruses. *Virology* 66, 616-619.

Lo K, Landau NR, Smale ST. *Mol. Cell. Biol.* 11:5229-5243 (1991)

Maniatis T, Fritsch EF and Sambrook J. (1982) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbour Laboratory, Cold Spring Harbour, NY.

Saiki RK, Gelfand DH, Stoffel S, Scharf SC, Higuchi R, Horn GT, Mullis KB and Erlich HA. (1987). Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239, 487-491.

Sambrook J Fritsch EF, and Maniatis T. (1989). *Molecular Cloning a Laboratory Manual*, 2nd ed. Cold Spring Harbour Laboratory, Cold Spring Harbour New York.

Smith GE, Summers MD and Fraser MJ. (1983). Production of human beta interferon in insect cells infected with a baculovirus vector. *Mol. Cell. Biol.*, 3, 2156-2165.

Stranstrom H, Verjalainen P, Meoning V Hunsmann G, Schwartz H. and Schafer W. (1974). C- type particles produced by a permanent cell line from a leukemic pig. 1 Origin and properties of host cells and some evidence for the occurence of C-Type like particles *Virology* 57, 175-178.

Todaro GJ, Benveniste RE, Lieber MM and Sherr CJ. (1974). Characterizaation of a type C virus released from the porcine cell line PK (15). *Virology* 58, 65-74.